AB-NK025-7 CDK1-X Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase CDK1 (CDC2)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Target Protein

Name Long:

Cyclin-dependent protein-serine kinase 1; Cell division control protein 2 homologue

Cdc2; CDC28; CDC28A; CDC2A; Cell division control protein 2; Cell division cycle 2, G1 to S and G2 to M; Cyclin-dependent kinase 1; P34 protein kinase; Kinase Cdc2; MPF; DKFZp686L20222; MGC111195; ENSG00000170312

UniProt ID:

P06493

Sequence Predicted Mass (KDa):

34.095 (297 AA; P06493); 27.503 (240 AA; P06493-2)

Observed SDS-PAGE Mass (KDa):

30-35

Immunogen

Antibody Immunogen Source: Human CDK1 (CDC2) sequence peptide

Antibody Immunogen Sequence: PLFHGDSEIDQLFRIFALGTPGGC

Location in Target: Corresponds to amino acid residues P202 to P223; In the protein kinase catalytic subdomain X region.

Peptide Type: For pan-specific recognition of target expression levels.

Target Phosphosite: Not phosphorylated

Production

Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Amount:	25 μg
Antibody Concentration:	1 mg/ml
Lot Number:	160520
Storage Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.

AB-NK025-7 CDK1-X Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications

Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	2 μg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Detection by Immunoblotting in Cell/Tissue Lysates:	Weak immunoreactivity of a target-sized protein by Western blotting in brain, testes and thymus.
Overall Antibody Specificity:	Medium selectivity
Antibody Cross Reactivities:	Very strong detection of 46 KDa protein in mouse brain and 38 KDa protein in mouse skeletal muscle that may be GSK3-beta, which was also detected in various human cancer cell lines (e.g. A431, HEK-293, HepG2, Jurkat and MCF7).

This product is for in vitro research use only and is not intended for use in humans or animals.