AB-NK048-7 PRKDC-4 (DNAPK) Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase DNAPK (PRKDC)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

	ΔT	$\mu \sim 1$	מומ
Targ			
	U .		

Name Long:	DNA-dependent protein kinase catalytic subunit
Alias:	DNA-PK catalytic subunit; DNA-PKcs; DNPK1; P460; PRKD; PRKDC; XRCC7; HYRC; HYRC1; p350; ENSG00000121031
UniProt ID:	P78527
Sequence Predicted Mass (KDa):	469.089 (4128 AA; P78527); 465.501 (4097 AA; P78527-2)
Observed SDS-PAGE Mass (KDa):	460-480

Immunogen

Antibody Immunogen Source:	Human DNAPK (PRKDC) sequence peptide Cat. No.: PE-01ARO85
Antibody Immunogen Sequence:	CLMDQATDPNILGRTW
Location in Target:	Corresponds to amino acid residues C4106 to W4121; C-terminus
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated

Production

Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Amount:	25 μg
Antibody Concentration:	0.75 mg/ml
Lot Number:	140724
Storage Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.

AB-NK048-7 PRKDC-4 (DNAPK) Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications

Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	2 μg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals

This product is for in vitro research use only and is not intended for use in humans or animals.