AB-NK052-6 EGFR-2 Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-tyrosine kinase EGFR (ErbB1)



Email: info@kinexus.ca Phone: 604-323-2547

Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Target Protein	
Name Long:	Epidermal growth factor receptor-tyrosine kinase
Alias:	EGFR; Epidermal growth factor receptor; ErbB-1; ErbB, mENA; HER1; Receptor tyrosine-protein kinase ErbB-1; V-erb-b oncogene homologue; PIG61; CCDS5514.1; ENSG00000146648
UniProt ID:	P00533
Sequence Predicted Mass (KDa):	134.277 (1210 AA; P00533); 77.312 (705 AA; P00533-3); 69.228 (628 AA; P00533-4); 44.664 (405 AA; P00533-2)
Observed SDS-PAGE Mass (KDa):	160-180
Immunogen	
Antibody Immunogen Source:	Human EGFR (ErbB1) sequence peptide Cat. No.: PE-01ATT90
Antibody Immunogen Sequence:	CVIQGDERMHLPSPT
Location in Target:	Corresponds to amino acid residues V980 to T993; Just after the catalytic domain of the kinase.
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated
Production Antibody Host Species:	Rabbit
Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH
	7.0 with saturated Tris.
Antibody Amount:	
Antibody Amount: Antibody Concentration:	7.0 with saturated Tris.
	7.0 with saturated Tris. 25 μg
Antibody Concentration:	7.0 with saturated Tris. 25 μg 1 mg/ml





Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications	
Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	2 µg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Antibody Positive Controls:	Weak reactivity with recombinant human EGFR on protein dot blots.
Overall Antibody Specificity:	Very high selectivity
Antibody Cross Reactivities:	No immunoreactivity on protein dot blots with recombinant human ErbB2 and ErbB4

This product is for in vitro research use only and is not intended for use in humans or animals.