AB-NK055-4 ERK1-NT (ERK1-4) Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase ERK1 (MAPK3)



Email: info@kinexus.ca Phone: 604-323-2547

Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Target Protein	
Name Long:	Extracellular regulated protein-serine kinase 1 (p44 MAP kinase)
Alias:	ERK-1; ERT2; Insulin-stimulated MAP2 kinase; Kinase ERK1; MAP kinase 1; MAPK 1; MAPK1; MAPK3; PRKM3; p44ERK1; p44MAPK; MGC20180; ENSG00000102882
UniProt ID:	P27361
Sequence Predicted Mass (KDa):	43.136 (379 AA; P27361); 40.088 (357 AA; P27361-3); 38.275 (335 AA; P27361-2)
Observed SDS-PAGE Mass (KDa):	42-47
Immunogen	
Antibody Immunogen Source:	Human ERK1 (MAPK3) sequence peptide Cat. No.: PE-01AQS95
Antibody Immunogen Sequence:	CGGGGGEPRRTEGVGPGVPGEVEMVKGGC
Location in Target:	Corresponds to amino acid residues G8 to K32; N-terminus
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated
Production Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by
Production Method:	mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
	mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH
Production Method: Antibody Amount: Antibody Concentration:	mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Amount:	mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Amount: Antibody Concentration:	mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. 25 μ g 0.5 / 0.16 mg/ml

AB-NK055-4 ERK1-NT (ERK1-4) Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications	
Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	1 µg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Detection by Immunoblotting in Cell/Tissue Lysates:	Very strong immunoreactivity of a target-sized protein by Western blotting in mouse brain, muscle and testes.
Overall Antibody Specificity:	Very high selectivity
Antibody Cross Reactivities:	No significant cross-reactive proteins detected in A431, HEK-293, HeLa, HepG2, Jurkat and MCF7 cells.

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit <u>www.kinexusproducts.ca</u> or contact us at 1-866-KINEXUS(546-3987)