AB-NK055-7 ERK1-I Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase ERK1 (MAPK3)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

-	4	_			
Tarq	Δt	Рr	\mathbf{C}	ים	ın
ıaıu	$-\iota$		U	U	

Name Long:	Extracellular regulated protein-serine kinase 1 (p44 MAP kinase)
Alias:	ERK-1; ERT2; Insulin-stimulated MAP2 kinase; Kinase ERK1; MAP kinase 1; MAPK 1; MAPK1; MAPK3; PRKM3; p44ERK1; p44MAPK; MGC20180; ENSG00000102882
UniProt ID:	P27361
Sequence Predicted Mass (KDa):	43.136 (379 AA; P27361); 40.088 (357 AA; P27361-3); 38.275 (335 AA; P27361-2)
Observed SDS-PAGE Mass (KDa):	42-47

Immunogen

Antibody Immunogen Source:	Human ERK1 (MAPK3) sequence peptide Cat. No.: PE-01AQH40
Antibody Immunogen Sequence:	GLAYIGEGAYGMVYKAGGC
Location in Target:	Corresponds to amino acid residues G38 to V50; Kinase catalytic Subdomain I
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated

Production

Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Amount:	25 μg
Antibody Concentration:	0.15 mg/ml
Lot Number:	160520
Storage Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.

AB-NK055-7 ERK1-I Antibody

Applications

Overall Antibody Specificity:

Antibody Cross Reactivities:



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

No significant cross-reactive proteins detected in A431, HepG2, MCF7 and HeLa

7 (ppiloations			
Product Use:	Western blotting Antibody microarrays		
Antibody Dilution Recommended:	4 μg/ml for immunoblotting		
Antibody Species Reactivity:	Human, mouse, rat and many other mammals		
Detection by Immunoblotting in	Strong immunoreactivity of a target-sized protein by Western blotting in mouse		
Cell/Tissue Lysates:	brain.		

cells, except for a weak 82 KDa protein.

High selectivity

This product is for in vitro research use only and is not intended for use in humans or animals.