AB-NK059-4

p38g-2 Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase p38g MAPK (MAPK12, ERK6)

Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3



Email: info@kinexus.ca Phone: 604-323-2547

	1 -		
loro	Ot L	roti	מור
Targ			
	O	100	2111

Name Long:	Mitogen-activated protein-serine kinase p38 gamma; Mitogen-activated protein kinase 12
Alias:	ERK3; ERK5; ERK6; MAPK12; MK12; p38 MAPK-gamma; PRKM12; SAPK3; Stress-activated protein kinase-3; CCDS14089.1; ENSG00000188130
UniProt ID:	P53778
Sequence Predicted Mass (KDa):	41.940 (367 AA; P53778); 40.808 (357 AA; P53778-2)
Observed SDS-PAGE Mass (KDa):	38-42

Immunogen

Antibody Immunogen Source:	Human p38g MAPK (MAPK12, ERK6) sequence peptide Cat. No.: PE-01AYK90	
Antibody Immunogen Sequence:	CVQKYDDSFDDVDRTL	
Location in Target:	Corresponds to amino acid residues V323 to L337; Post-kinase last alpha-chain	
Peptide Type:	For pan-specific recognition of target expression levels.	
Target Phosphosite:	Not phosphorylated	

Production

ody Host Species: ody Type:	Rabbit Polyclonal		
ody Type:	Polyclonal		
7J . J P C .	Polyclonal		
ody lg Isotype Clone Lot:	Immunoglobulin G		
ction Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.		
ody Amount:	25 μg		
ody Concentration:	1 mg/ml		
umber:	140724		
ge Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol		
ge Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.		

AB-NK059-4 p38g-2 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Αp	pli	cati	ions

7 (ppiloations			
Product Use:	Western blotting Antibody microarrays		
Antibody Dilution Recommended:	1 μg/ml for immunoblotting		
Antibody Species Reactivity:	Human, mouse, rat and many other mammals		
Antibody Positive Controls: Medium-strong immunoreactivity with recombinant human p38g on protein do blots.			
Overall Antibody Specificity:	Very high selectivity		
Antibody Cross Reactivities:	Medium-strong with recombinant human p38b, very weak immunoreactivity with p38d, and no immunoreactivity with p38a.		

This product is for in vitro research use only and is not intended for use in humans or animals.