AB-NK074-3 IGF1R-CT Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-tyrosine kinase IGF1R



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Target Protein

Name Long:

Insulin-like growth factor 1 receptor protein-tyrosine kinase

CD221; IGFIR; Insulin-like growth factor I receptor; JTK13; MGC142170; MGC142172

UniProt ID: P08069

Sequence Predicted Mass (KDa): 154.793 (1367 AA; P08069)

Observed SDS-PAGE Mass (KDa): 90-100, 130

Immunogen

Antibody Immunogen Source: Human IGF1R sequence peptide Cat. No.: PE-01BDF95

Antibody Immunogen Sequence: HMNGGRKNERALPLPQSSTC

Location in Target: Corresponds to amino acid residues H1348 to C1367; C-terminus

Peptide Type: For pan-specific recognition of target expression levels.

Target Phosphosite: Not phosphorylated

Production

Rabbit
Polyclonal
Immunoglobulin G
The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
0.5 mg/ml
160807
Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol
For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.

AB-NK074-3 IGF1R-CT Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications

Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	2 μg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals

This product is for in vitro research use only and is not intended for use in humans or animals.