AB-NK084-3 JAK1-1 Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-tyrosine kinase JAK1



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email:	info@kinexus.ca
Phone	: 604-323-2547

Target Protein	
Name Long:	Janus protein-tyrosine kinase 1
Alias:	JAK1A; Janus kinase 1; JAK1B; CE2.7.10.2; JAK-1; JTK3; JAK1_ENST00000342505; CCDS41346; Q0VGA4; Q4LDX3; Q6P669; Q8IV82; ENSG00000162434
UniProt ID:	P23458
Sequence Predicted Mass (KDa):	133.277 (1154 AA; P23458)
Observed SDS-PAGE Mass (KDa):	125-135
Immunogen	
Antibody Immunogen Source:	Human JAK1 sequence peptide Cat. No.: PE-01AUM99
Antibody Immunogen Sequence:	CFFRAIMRDINKLEEQ
Location in Target:	Corresponds to amino acid residues F837 to Q851; Kinase last alpha-chain
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated
Production	
Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The
	sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Amount:	immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH
Antibody Amount: Lot Number:	immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
-	immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. 25 μg





Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications	
Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	1 µg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Overall Antibody Specificity:	Low selectivity
Antibody Cross Reactivities:	No immunoreactivity on protein dot blots with recombinant human JAK2, JAK3 and Tyk2.

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit <u>www.kinexusproducts.ca</u> or contact us at 1-866-KINEXUS(546-3987)