AB-NK103-4 MKK4-1 Antibody

Pan-specific polyclonal antibody for monitoring the expression of human dual specificity protein kinase MKK4 (MAP2K4, MEK4)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Target Protein	
Name Long:	MAPK/ERK protein-serine kinase 4 (MKK4); Dual specificity mitogen-activated protein kinase kinase 4
Alias:	c-Jun N- terminal kinase kinase 1; C-JUN N-terminal kinase kinase 1; JNK activating kinase 1; JNK kinase 1; JNKK1; Kinase SEK1; MAP2K4; SEK1; SERK1; SAPKK1; SAPKK-1; SKK1; CCDS11162.1; ENSG00000065559
UniProt ID:	P45985
Sequence Predicted Mass (KDa):	45.584 (410 AA; P45985-2); 44.288 (399 AA; P45985)
Observed SDS-PAGE Mass (KDa):	42-46
Immunogen	
Antibody Immunogen Source:	Human MKK4 (MAP2K4, MEK4) sequence peptide Cat. No.: PE-01AWE99
Antibody Immunogen Sequence:	CGKLKISPEQHWDFTA
Location in Target:	Corresponds to amino acid residues G85 to A99;
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated
Production Antibody Host Species:	Rabbit
	Rabbit Polyclonal
Antibody Host Species:	
Antibody Host Species: Antibody Type:	Polyclonal
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot:	PolyclonalImmunoglobulin GThe immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method:	PolyclonalImmunoglobulin GThe immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: Antibody Amount:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. 25 μg
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: Antibody Amount: Antibody Concentration:	PolyclonalImmunoglobulin GThe immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.25 μg1 mg/ml





Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications	
Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	1 µg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Antibody Positive Controls:	Strong immunoreactivity with recombinant human MKK4 on protein dot blots.
Overall Antibody Specificity:	Low selectivity
Antibody Cross Reactivities:	No immunoreactivity on protein dot blots with recombinant human MEK1, MEK2 and MKK6.

This product is for in vitro research use only and is not intended for use in humans or animals.