AB-NK104-2 MEK5-PNT (MEK5-4) Antibody

Pan-specific polyclonal antibody for monitoring the expression of human dual specificity protein kinase MEK5 (MAP2K5, MKK5)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

| Targ | Δt | uro | tain |
|------|------------|-----|------|
| | | _ | |
| | | | |

Name Long: MAPK/ERK protein-serine kinase 5 (MKK5); Dual specificity mitogen-activated protein kinase kinase 5

OVA4. LETAZAFA: MAD kingga kingga F: MADOKE: MADK EDK

CXA1; HsT17454; MAP kinase kinase 5; MAP2K5; MAPK,ERK kinase 5; MAPKK5; MKK5; HsT17454; MAPK/ERK kinase 5; MAPKK5; MKK5; HsT17454;

ENSG00000137764

UniProt ID: Q13163

Sequence Predicted Mass (KDa): 50.112 (448 AA; Q13163); 49.497 (444 AA; Q13163-3); 48.968 (438 AA;

Q13163-2); 46.131 (412 AA; Q13163-4)

Observed SDS-PAGE Mass (KDa): 47-52

Immunogen

Antibody Immunogen Source: Human MEK5 (MAP2K5, MKK5) sequence peptide

Antibody Immunogen Sequence: AFEYEDEDGDRITVRSC

Location in Target: Corresponds to amino acid residues A59 to S74; Post-N-terminus

Peptide Type: For pan-specific recognition of target expression levels.

Target Phosphosite: Not phosphorylated

Production

Production Method:

Antibody Host Species: Rabbit
Antibody Type: Polyclonal

Antibody Ig Isotype Clone Lot: Immunoglobulin G

The immunizing peptide was produced on an A.B.I. automated peptide synthesizer. The peptide was cleaved from the synthesis resin with HF and purified by reverse-phase hplc chromatography. Purity was assessed by

purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by amino acid analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand

White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution

was neutralized to pH 7.0 with saturated Tris.

Antibody Amount: 25 µg

Antibody Concentration: 0.25 mg/ml

Lot Number: 10320

Storage Buffer: 100 mM Tris-glycine, pH 7.0

For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months, but either 0.1% sodium azide or 0.05%

Thimerasol should be added. Avoid repeated freeze-thaw cycles.

AB-NK104-2 MEK5-PNT (MEK5-4) Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications

| Product Use: | Western blotting Antibody microarrays | |
|---------------------------------------|--|--|
| Antibody Dilution Recommended: | 2 μg/ml for immunoblotting | |
| Antibody Species Reactivity: | Human, mouse, rat and many other mammals | |

This product is for in vitro research use only and is not intended for use in humans or animals.