AB-NK121-4

p38d-3 Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase p38d MAPK (MAPK13)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

37-45

Email: info@kinexus.ca Phone: 604-323-2547

	чат	Prof	
1011			

Name Long:	Mitogen-activated protein-serine kinase p38 delta; Mitogen-activated protein kinase 13
Alias:	MAP kinase p38 delta; MAPK13; Mitogen-activated protein kinase p38 delta; MK13; P38 MAPK-delta;p38delta; PRKM13; Stress-activated protein kinase-4; SAPK4; p38d MAPK; ENSG00000156711
UniProt ID:	O15264
Sequence Predicted Mass (KDa):	42.090 (365 AA; O15264); 28.779 (257 AA; O15264-2)

Immunogen

Observed SDS-PAGE Mass (KDa):

Antibody Immunogen Source:	Human p38d MAPK (MAPK13) sequence peptide Cat. No.: PE-01AYI90	
Antibody Immunogen Sequence:	CPIARKDSRRRSGMKL	
Location in Target:	Corresponds to amino acid residues P351 to L365; C-terminus	
Peptide Type:	For pan-specific recognition of target expression levels.	
Target Phosphosite:	Not phosphorylated	

Production

Antibody Host Species:	Rabbit	
Antibody Type:	Polyclonal	
Antibody Ig Isotype Clone Lot:	Immunoglobulin G	
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.	
Antibody Amount:	25 μg	
Antibody Concentration:	0.75 mg/ml	
Lot Number:	140724	
Storage Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol	
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.	

AB-NK121-4 p38d-3 Antibody

Applications

Antibody Cross Reactivities:



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications		
Product Use:	Western blotting Antibody microarrays	
Antibody Dilution Recommended:	1 μg/ml for immunoblotting	
Antibody Species Reactivity:	Human, mouse, rat and many other mammals	
Antibody Positive Controls:	Medium immunoreactivity with recombinant human p38d on protein dot blots.	
Overall Antibody Specificity:	High selectivity	
Antibody Cross Posstivities	Weak immunoreactivity on protein dot blots with recombinant human p38g and	

p38b, and no immunoreactivity with p38a.

This product is for in vitro research use only and is not intended for use in humans or animals.