AB-NK122-2 PAK1-NT (PAK1-1) Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase PAK1 (PAKa)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

60-67

Email: info@kinexus.ca Phone: 604-323-2547

11	$r \cap t$	'AID
	et P	et Prot

Name Long:

p21-activated kinase 1 alpha; Protein-serine/threonine kinase PAK 1

ADRB2; Alpha-PAK; CDC42,RAC effector kinase PAK-A; CDC42/RAC effector kinase PAK-A; Kinase PAK-A; Kinase PAK-A; MUK2; P21 protein (Cdc42/Rac)-activated kinase 1; P65-PAK; STE20, yeast; P68-PAK; PAK 1; PAK-1; PAKA; Protein kinase MUK2; PAKalpha; MGC130000; MGC130001; CCDS8250.1; Q13153; ENSG00000149269

UniProt ID:

Q13153

Sequence Predicted Mass (KDa):

61.632 (553 AA; Q13153-2); 60.647 (545 AA; Q13153)

Immunogen

Observed SDS-PAGE Mass (KDa):

Antibody Immunogen Source: Human PAK1 (PAKa) sequence peptide

Antibody Immunogen Sequence: MSNNGLDVQDKPC

Location in Target: Corresponds to amino acid residues M1 to P12; N-terminus

Peptide Type: For pan-specific recognition of target expression levels.

Target Phosphosite: Not phosphorylated

Production

Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced on an A.B.I. automated peptide synthesizer. The peptide was cleaved from the synthesis resin with HF and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by amino acid analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Amount:	25 μg
Antibody Concentration:	0.25 mg/ml
Lot Number:	10320
Storage Buffer:	100 mM Tris-glycine, pH 7.0
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months, but either 0.1% sodium azide or 0.05% Thimerasol should be added. Avoid repeated freeze-thaw cycles.

AB-NK122-2 PAK1-NT (PAK1-1) Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications

Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	2 μg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Antibody Cross Reactivities:	Cross-reactivity with 60 KDa and 90 KDa proteins in various rat tissues.

This product is for in vitro research use only and is not intended for use in humans or animals.