AB-NK155-7 Raf1-3 Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase Raf1 (RafC)



Email: info@kinexus.ca Phone: 604-323-2547

Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Target Protein	
Name Long:	Raf1 proto-oncogene-encoded protein-serine kinase
Alias:	cRAF; c-RAF; c-Raf; Kinase Raf1; Murine leukemia viral (V-raf-1) oncogene homologue 1; NS5; Protein kinase raf 1; Raf proto-oncogene serine/threonine- protein kinase; RAF1; V-raf-1 murine leukemia viral oncogene 1; CCDS2612.1; ENSG00000132155; B4E1N6; Q59HC8
UniProt ID:	P04049
Sequence Predicted Mass (KDa):	75.395 (668 AA; P04049-2); 73.052 (648 AA; P04049)
Observed SDS-PAGE Mass (KDa):	70-75
Immunogen	
Antibody Immunogen Source:	Human Raf1 (RafC) sequence peptide Cat. No.: PE-01AYO99
Antibody Immunogen Sequence:	CLIGEELQVDFLDHVP
Location in Target:	Corresponds to amino acid residues L121 to P135
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated
Production	
	Rabbit
Antibody Host Species:	Rabbit
Antibody Host Species: Antibody Type:	Polyclonal
Antibody Host Species:	
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: Antibody Amount:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. 25 μg
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: Antibody Amount: Antibody Concentration:	PolyclonalImmunoglobulin GThe immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.25 μg1 mg/ml





Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications		
Product Use:	Western blotting Antibody microarrays	
Antibody Dilution Recommended:	2 µg/ml for immunoblotting	
Antibody Species Reactivity:	Human, mouse, rat and many other mammals	
Overall Antibody Specificity:	High selectivity	
Antibody Cross Reactivities:	No immunoreactivity on protein dot blots with recombinant human B-Raf.	

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit <u>www.kinexusproducts.ca</u> or contact us at 1-866-KINEXUS(546-3987)