AB-NK202 PKG1a-NT (PKG1a-1) Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase PKG1a (PRKG1A)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Target Protein

Name Long:	cGMP-dependent protein kinase 1, alpha isozyme
Alias:	cGK 1 alpha; cGKI-alpha; cGKI-beta; cGMP-dependent protein kinase 1, alpha isozyme; DKFZp686K042; FLJ36117; KGP1A; Kinase PKG1; PGK; PKG; PKG1-alpha; Protein kinase, cGMP-dependent, type I
UniProt ID:	Q13976
Sequence Predicted Mass (KDa):	77.804 (686 AA; Q13976-2); 76.364 (671 AA; Q13976); 44.185 (389 AA; Q13976-3)
Observed SDS-PAGE Mass (KDa):	77-83

Immunogen

Antibody Immunogen Source:	Human PKG1a (PRKG1A) sequence peptide
Antibody Immunogen Sequence:	ELEEDFAKILMLKEEC
Location in Target:	Corresponds to amino acid residues E3 to E17; N-terminus
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated

Production

Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced on an A.B.I. automated peptide synthesizer. The peptide was cleaved from the synthesis resin with HF and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by amino acid analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Amount:	25 μg
Antibody Concentration:	0.5 mg/ml
Lot Number:	10320
Storage Buffer:	100 mM Tris-glycine, pH 7.0
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months, but either 0.1% sodium azide or 0.05% Thimerasol should be added. Avoid repeated freeze-thaw cycles.

AB-NK202 PKG1a-NT (PKG1a-1) Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications

Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	2 μg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals

This product is for in vitro research use only and is not intended for use in humans or animals.