AB-NK243-1 PDGFRB-2 Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-tyrosine kinase PDGFRb



Email: info@kinexus.ca

Phone: 604-323-2547

Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Target Protein	
Name Long:	Platelet-derived growth factor receptor kinase beta
Alias:	CD140b; JTK12; Kinase PDGFR-beta; PDGFR; PDGFR1; PDGFRb; PDGFR- beta; PDGF-R-beta; Platelet-derived growth factor receptor, beta polypeptide; BC032224; J03278; M21616; CCDS4303.1; ENSG00000113721
UniProt ID:	P09619
Sequence Predicted Mass (KDa):	123.968 (1106 AA; P09619); 37.412 (336 AA; P09619-2)
Observed SDS-PAGE Mass (KDa):	115-130
Immunogen	
Antibody Immunogen Source:	Human PDGFRb sequence peptide Cat. No.: PE-01AUY99
Antibody Immunogen Sequence:	CKGDVALPVPYDHQRG
Location in Target:	Corresponds to amino acid residues K164 to G178;
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated
Production	
Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
	i olyolonai
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Antibody Ig Isotype Clone Lot: Production Method:	-
	Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH
Production Method:	Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Production Method: Antibody Amount:	Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. 25 µg
Production Method: Antibody Amount: Antibody Concentration:	Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. 25 μg 1 mg/ml

For long term storage, keep frozen at -40°C or lower. Stock solution can be kept

at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.

Storage Conditions and Stability:

AB-NK243-1 PDGFRB-2 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications	
Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	2 µg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Antibody Positive Controls:	Medium-strong immunoreactivity with recombinant human PDGFRA on protein dot blots.
Overall Antibody Specificity:	High selectivity
Antibody Cross Reactivities:	No immunoreactivity on protein dot blots with recombinant human PDGFRA.

This product is for in vitro research use only and is not intended for use in humans or animals.