AB-NK247-3 Obscn-3 Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase Obscn



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Target Protein	
Name Long:	Obscurin-myosin light chain kinase
Alias:	Obscurin; UNC89; ARHGEF30; Q96AA2; ENSG00000154358
UniProt ID:	Q5VST9
Sequence Predicted Mass (KDa):	924.971 (8483 AA; Q5VST9-6); 868.571 (7969 AA; Q5VST9-2); 868.484 (7968 AA; Q5VST9); 721.547 (6620 AA; Q5VST9-3); 427.046 (3911 AA; Q5VST9-5)
Observed SDS-PAGE Mass (KDa):	36.6-45
Immunogen	
Antibody Immunogen Source:	Human Obscn sequence peptide Cat. No.: PE-01AXZ99
Antibody Immunogen Sequence:	CRRALLYKRHNLAQVR
Location in Target:	Corresponds to amino acid residues R7955 to R7968; C-terminus
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated
Production	
Production	
Antibody Host Species:	Rabbit
	Rabbit Polyclonal
Antibody Host Species:	
Antibody Host Species: Antibody Type:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: Antibody Amount:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. 25 μg
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.

Storage Buffer:Phosphate buffered saline (PBS) pH7.4, 0.05% ThimerasolStorage Conditions and Stability:For long term storage, keep frozen at -40°C or lower. Stock solution can be kept

Storage Conditions and Stability: at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.





Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Product Use: Western blotting Antibody microarrays Antibody Dilution Recommended 2 ug/ml for immunobletting	
Antibody Dilution Decommonded	
Antibody Dilution Recommended: 2 µg/ml for immunoblotting	
Antibody Species Reactivity: Human, mouse, rat and many other mammals	

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit <u>www.kinexusproducts.ca</u> or contact us at 1-866-KINEXUS(546-3987)