## AB-NK255-1 WNK4-1 Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase WNK4 (PRKWNK4)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Target Protein	
Name Long:	Serine/threonine-protein kinase WNK4
Alias:	Kinase WNK4; PRKWNK4; Protein kinase with no lysine 4; Protein kinase, lysine-deficient 4; WNK lysine deficient protein kinase 4; PHA2B; CCDS11439.1; ENSG00000126562
UniProt ID:	Q96J92
Sequence Predicted Mass (KDa):	134.739 (1243 AA; Q96J92); 126.577 (1165 AA; Q96J92-3); 73.370 (663 AA; Q96J92-2)
Observed SDS-PAGE Mass (KDa):	110, 140-180
Immunogen	
Antibody Immunogen Source:	Human WNK4 (PRKWNK4) sequence peptide Cat. No.: PE-01AZH85
Antibody Immunogen Sequence:	MLASPATETTVLMSQC
Location in Target:	Corresponds to amino acid residues M1 to Q15; N-terminus
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated
Production	
	Rabbit
Antibody Host Species:	
Antibody Host Species: Antibody Type:	Polyclonal
Antibody Host Species:	
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: Antibody Amount:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. 25 μg
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: Antibody Amount: Antibody Concentration:	PolyclonalImmunoglobulin GThe immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.25 μg1 mg/ml





Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications	
Product Use:	Western blotting   Antibody microarrays
Antibody Dilution Recommended:	2 µg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
<b>Overall Antibody Specificity:</b>	Medium selectivity
Antibody Cross Reactivities:	No immunoreactivity on protein dot blots with recombinant human WNK1.

This product is for in vitro research use only and is not intended for use in humans or animals.