AB-NK300-1 HGK-NT Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase HGK (ZC1)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

	4		1	
arc	1 O T	$r \cap$	tΔ	ın
arc	ıCı	ı	เบ	ш

3			
Name Long:	Mitogen-activated protein kinase kinase kinase 4		
Alias:	FLH21957; Hepatocyte progenitor kinase-like/germinal center kinase-like kinase; HGK; HPK/GCK-like kinase HGK; M4K4; MAP4K4; MAPK kinase kinase kinase 4; MAPK/ERK kinase kinase kinase 4; MEK kinase kinase 4; MEKKK 4; MEKKK4; Mitogen-activated protein kinase kinase kinase kinase 4; Nck interacting kinase; NIK; FLJ10410; FLJ20373; FLJ90111; KIAA0687; NIK; O95819; ENSG0000007105		
UniProt ID:	O95819		
Sequence Predicted Mass (KDa):	150.913 (1319 AA; O95819-3); 142.101 (1239 AA; O95819); 141.929 (1242 AA; O95819-5); 138.418 (1211 AA; O95819-2); 133.401 (1165 AA; O95819-4)		
Observed SDS-PAGE Mass (KDa):	135-150		

Immunogen

Antibody Immunogen Source:	Human HGK (ZC1) sequence peptide Cat. No.: PE-01BFM99T
Antibody Immunogen Sequence:	ANDSPAKSLVDIDLSSLRDC
Location in Target:	Corresponds to amino acid residues A2 to D20; N-terminus
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated

Production

Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Antibody Amount:	25 μg
Antibody Concentration:	1 mg/ml
Lot Number:	160630
Storage Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.

AB-NK300-1 HGK-NT Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications		
Product Use:	Western blotting Antibody microarrays	
Antibody Dilution Recommended:	2 μg/ml for immunoblotting	
Antibody Species Reactivity:	Human, mouse, rat and many other mammals	
Overall Antibody Specificity:	High selectivity	
Antibody Cross Reactivities:	No significant cross-reactivities detected in various human cancer cell lines, including A431, HEK-293, HeLa, HepG2, Jurkat and MCF7 cells, except for very strong ~82 KDa and ~25-26 KDa cross-reactive proteins. No cross-reactivities detected in mouse tissues, except for very strong ~85 and ~90 KDa proteins in mouse brain, skeletal muscle and testes.	