AB-NN178-1 PDK4 Antibody

Pan-specific polyclonal antibody for monitoring the expression of human protein-serine/threonine kinase PDK4 (PDHK4)



Email: info@kinexus.ca

Phone: 604-323-2547

Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Target Protein	
Name Long:	Pyruvate dehydrogenase kinase isoform 4; [Pyruvate dehydrogenase
Alias:	[lipoamide]] kinase isozyme 4, mitochondrial Pyruvate dehydrogenase kinase isoform 4; PDK4;
UniProt ID:	Q16654
Sequence Predicted Mass (KDa):	46.469 (411 AA; Q16654)
Observed SDS-PAGE Mass (KDa):	45-55
Immunogen	
Antibody Immunogen Source:	Human PDK4 (PDHK4) sequence peptide
Antibody Immunogen Sequence:	
Location in Target:	Corresponds to amino acid residues to ;
Peptide Type:	For pan-specific recognition of target expression levels.
Target Phosphosite:	Not phosphorylated
Draduction	
Production	
Antibody Host Species:	Rabbit
	Rabbit Polyclonal
Antibody Host Species:	
Antibody Host Species: Antibody Type:	Polyclonal
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.





Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications		
Product Use:	Western blotting Antibody microarrays	
Antibody Dilution Recommended:	2 µg/ml for immunoblotting	
Antibody Species Reactivity:	Human, mouse, rat and many other mammals	

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit <u>www.kinexusproducts.ca</u> or contact us at 1-866-KINEXUS(546-3987)