AB-PK628 FAK-pY576+pY577 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human protein-tyrosine kinase FAK (PTK2)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Target Protein	
Name Long:	Focal adhesion protein-tyrosine kinase
Alias:	FADK1; FAK1; pp125FAK; FRNK; Protein-tyrosine kinase 2; PTK2; CCDS6381.1; ENSG00000169398
UniProt ID:	Q05397
Sequence Predicted Mass (KDa):	120.900 (1065 AA; Q05397-5); 119.233 (1052 AA; Q05397); 114.253 (1006 AA; Q05397-7); 99.358 (879 AA; Q05397-2); 63.201 (554 AA; Q05397-3); 48.967 (431 AA; Q05397-4); 39.874 (360 AA; Q05397-6)
Observed SDS-PAGE Mass (KDa):	105-115
Immunogen	
Antibody Immunogen Source:	Human FAK (PTK2) sequence peptide Cat. No.: PE-04AGR90
Antibody Immunogen Sequence:	DST(pY)(pY)KAS(bA)C (bA) = beta-alanine
Location in Target:	Corresponds to amino acid residues D573 to S580; In protein kinase catalytic domain activation T-loop between subdomains VII and VIII.
Peptide Type:	For phosphosite-specific recognition of target.
Target Phosphosite:	Tyr-576+Tyr-577
Production Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH
	7.0 with saturated Tris. This antibody was also subject to negative purification over phosphotyrosine-agarose.
Antibody Amount:	
Antibody Amount: Antibody Concentration:	over phosphotyrosine-agarose.
	over phosphotyrosine-agarose. 25 μg
Antibody Concentration:	over phosphotyrosine-agarose. 25 µg 1 mg/ml

AB-PK628 FAK-pY576+pY577 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications	
Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	2 μg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other vertebrates; Phosphosite is highly conserved in diverse species
Antibody Positive Controls:	Medium immunoreactivity with immunogen peptide on dot blots.
Overall Antibody Specificity:	High-very high selectivity
Antibody Cross Reactivities:	No significant cross-reactive proteins detected in A431 and MCF7 cells.

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit <u>www.kinexusproducts.ca</u> or contact us at 1-866-KINEXUS(546-3987)