# AB-PK666 ITK-pY512 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human protein-tyrosine kinase ITK



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

### Target Protein

Name Long:	Tyrosine-protein kinase ITK/TSK
Alias:	EMT; Lyk; PSCTK2; MGC126257; MGC126258; TLK; TSK; CCDS4336.1; ENSG00000113263
UniProt ID:	Q08881
Sequence Predicted Mass (KDa):	71.831 (620 AA; Q08881)
Observed SDS-PAGE Mass (KDa):	68-72

### Immunogen

Antibody Immunogen Source:	Human ITK sequence peptide Cat. No.: PE-04ANB90
Antibody Immunogen Sequence:	DDQ(pY)TSS(bA)C (bA) = beta-alanine
Location in Target:	Corresponds to amino acid residues D509 to S515; In protein kinase catalytic domain activation T-loop between subdomains VII and VIII.
Peptide Type:	For phosphosite-specific recognition of target.
Target Phosphosite:	Tyr-512

#### Production

Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification over phosphotyrosine-agarose.
Antibody Amount:	25 µg
Antibody Concentration:	1 mg/ml
Lot Number:	150305
Storage Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.

## AB-PK666 ITK-pY512 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

#### Applications

Product Use:	Western blotting   Antibody microarrays
Antibody Dilution Recommended:	2 μg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Antibody Positive Controls:	Very strong immunoreactivity with immunogen peptide on dot blots.
Detection by Immunoblotting in Cell/Tissue Lysates:	Weak immunoreactivity of a target-sized protein by Western blotting in Jurkat cells.
Overall Antibody Specificity:	High selectivity
Antibody Cross Reactivities:	No significant cross-reactivities detected in Jurkat cells, except phenylarsine oxide (PAO) increases 15 KDa protein immunoreactivity.

This product is for in vitro research use only and is not intended for use in humans or animals.