## AB-PK670 JNK1-pY185 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human protein-serine/threonine kinase JNK1 (MAPK8)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Target Protein	
Name Long:	Jun N-terminus protein-serine kinase (stress-activated protein kinase (SAPK)) 1
Alias:	c-Jun N-terminal kinase 1; JNK; JNK1A2; JNK1-alpha-2; JNK-46; JNK21B1/2; JUN N-terminal kinase; Kinase JNK1; MAPK8; MK08; PRKM8; SAPK1; SAPK1c; Stress-activated protein kinase JNK1; CCDS7225.1; ENSG00000107643
UniProt ID:	P45983
Sequence Predicted Mass (KDa):	48.296 (427 AA; P45983); 48.088 (427 AA; P45983-4); 44.229 (384 AA; P45983-2); 44.022 (384 AA; P45983-3); 35.333 (308 AA; P45983-5)
Observed SDS-PAGE Mass (KDa):	43-46
Immunogen	
Antibody Immunogen Source:	Human JNK1 (MAPK8) sequence peptide Cat. No.: PE-04AEM95
Antibody Immunogen Sequence:	MTP(pY)VVT(bA)C (bA) = beta-alanine
Location in Target:	Corresponds to amino acid residues M182 to T188; In protein kinase catalytic domain activation T-loop between subdomains VII and VIII.
Peptide Type:	For phosphosite-specific recognition of target.
Target Phosphosite:	Tyr-185
Production   Antibody Host Species:	Rabbit
Antibody Host Species: Antibody Type:	Polyclonal
Antibody Host Species:	Polyclonal Immunoglobulin G
Antibody Host Species: Antibody Type:	Polyclonal
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. This antibody was also subject to negative purification over phosphotyrosine-agarose.
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: Antibody Amount:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification over phosphotyrosine-agarose. 25 μg
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: Antibody Amount: Antibody Concentration:	PolyclonalImmunoglobulin GThe immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. This antibody was also subject to negative purification over phosphotyrosine-agarose.25 μg0.75 mg/ml

## AB-PK670 JNK1-pY185 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications	
Product Use:	Western blotting   Antibody microarrays
Antibody Dilution Recommended:	2 µg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Antibody Positive Controls:	Very strong immunoreactivity with immunogen peptide on dot blots. Medium- strong immunoreactivity with recombinant human JNK1 on protein dot blots.
Overall Antibody Specificity:	Very high selectivity
Antibody Cross Reactivities:	Strong immunoreactivity on protein dot blots with recombinant human JNK2, but very weak immunoreactivity with JNK3. No significant cross-reactive proteins detected in A431, A549, HeLa and MCF7 cells and in sea star oocytes, except for a ~77 KDa cross-reactive protein in A431 and Hela cells.

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit <u>www.kinexusproducts.ca</u> or contact us at 1-866-KINEXUS(546-3987)