AB-PK680 Lck-pY263+pY264 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human protein-tyrosine kinase Lck

KINEXUS

Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Target Protein

| Name Long: | Lymphocyte-specific protein-tyrosine kinase; Proto-oncogene tyrosine-protein kinase LCK |
|--------------------------------|---|
| Alias: | LSK; LSK-T; p56-LCK; pp58Lck; RP4-675E8.4; LSK (T-cell specific protein-tyrosine kinase; lck tyrosine kinase; membrane associated protein tyrosine kinase; proto-oncogene LCK; put.ptk; YT16; ENSG00000182866 |
| UniProt ID: | P06239 |
| Sequence Predicted Mass (KDa): | 61.190 (539 AA; P06239-3); 58.001 (509 AA; P06239); 40.866 (363 AA; P06239-2) |
| Observed SDS-PAGE Mass (KDa): | 50-58 |

Immunogen

| Antibody Immunogen Source: | Human Lck sequence peptide Cat. No.: PE-04ARP55 |
|------------------------------|--|
| Antibody Immunogen Sequence: | GEVWMG(pY)(pY)NGHT(bA)C (bA) = beta-alanine |
| Location in Target: | Corresponds to amino acid residues G257 to T268; In the kinase catalytic domain between subdomains I and II. |
| Peptide Type: | For phosphosite-specific recognition of target. |
| Target Phosphosite: | Tyr-263+Tyr-264 |

Production

| Troduction | |
|-----------------------------------|---|
| Antibody Host Species: | Rabbit |
| Antibody Type: | Polyclonal |
| Antibody Ig Isotype Clone Lot: | Immunoglobulin G |
| Production Method: | The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification over phosphotyrosine-agarose. |
| Antibody Amount: | 25 μg |
| Antibody Concentration: | 0.5 mg/ml |
| Lot Number: | 141003 |
| Storage Buffer: | Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol |
| Storage Conditions and Stability: | For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles. |

AB-PK680 Lck-pY263+pY264 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications

| Product Use: | Western blotting Antibody microarrays |
|--------------------------------|--|
| Antibody Dilution Recommended: | 2 μg/ml for immunoblotting |
| Antibody Species Reactivity: | Human, mouse, rat and many other mammals |

This product is for in vitro research use only and is not intended for use in humans or animals.