AB-PK725 MSK2-pT687 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human protein-serine/threonine kinase MSK2 (RPS6KA4)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

| Target Protein | |
|---|--|
| Name Long: | Ribosomal protein S6 kinase alpha-4 |
| Alias: | Kinase MSK2; KS6A4; Ribosomal protein kinase B; Ribosomal protein S6 kinase, 90kDa, polypeptide 4; RPS6KA4; RSK-B; Similar to ribosomal protein S6 kinase, 90kD, polypeptide 4; Similar to ribosomal protein S6 kinase, polypeptide 4; CCDS8073.1; ENSG00000162302 |
| UniProt ID: | O75676 |
| Sequence Predicted Mass (KDa): | 85.606 (772 AA; O75676); 84.884 (766 AA; O75676-2) |
| Observed SDS-PAGE Mass (KDa): | 83-90 |
| | |
| Immunogen | |
| Antibody Immunogen Source: | Human MSK2 (RPS6KA4) sequence peptide Cat. No.: PE-04AKR95 |
| Antibody Immunogen Sequence: | PLR(pT)PDC(bA)C (bA) = beta-alanine |
| Location in Target: | Corresponds to amino acid residues P684 to C690; In the C-terminal part of the kinase after the second catalytic domain |
| Peptide Type: | For phosphosite-specific recognition of target. |
| Target Phosphosite: | Thr-687 |
| | |
| Production Antibody Host Species: | Rabbit |
| | Rabbit Polyclonal |
| Antibody Host Species: | |
| Antibody Host Species: Antibody Type: | Polyclonal |
| Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: | Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification |
| Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: | Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification over phosphotyrosine-agarose. |
| Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: Antibody Amount: | Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification over phosphotyrosine-agarose. 25 μg |
| Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: Antibody Amount: Antibody Concentration: | PolyclonalImmunoglobulin GThe immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. This antibody was also subject to negative purification over phosphotyrosine-agarose.25 μg0.75 mg/ml |

AB-PK725 MSK2-pT687 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

| Applications | |
|--------------------------------|---|
| Product Use: | Western blotting Antibody microarrays |
| Antibody Dilution Recommended: | 2 µg/ml for immunoblotting |
| Antibody Species Reactivity: | Human, mouse, rat and many other mammals |
| Antibody Positive Controls: | Very strong immunoreactivity with immunogen peptide on dot blots. |
| Overall Antibody Specificity: | Very high selectivity |
| Antibody Cross Reactivities: | Almost no significant cross-reactivities in A431, Jurkat and T98G cells; in Jurkat cells, phenylarsine oxide (PAO) increases detection of 15 KDa protein. |

This product is for in vitro research use only and is not intended for use in humans or animals.