

# AB-PK764

## PKCa-pY195 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human protein-serine/threonine kinase PKCa (PRKCA)



**KINEXUS**

Address: 8755 Ash Street, Suite 1  
Vancouver, British Columbia,  
Canada V6P 6T3

Email: [info@kinexus.ca](mailto:info@kinexus.ca)  
Phone: 604-323-2547

### Target Protein

<b>Name Long:</b>	Protein-serine kinase C alpha
<b>Alias:</b>	AAG6; Kinase PKC-alpha; KPCA; PKC III; PKC-A; PKC-alpha; PKC-III; PRKACA; Protein kinase C, alpha; Protein kinase C, alpha type; MGC129900; MGC129901; CCDS11664.1; P17252; ENSG00000154229
<b>UniProt ID:</b>	P17252
<b>Sequence Predicted Mass (KDa):</b>	76.750 (672 AA; P17252)
<b>Observed SDS-PAGE Mass (KDa):</b>	75-90

### Immunogen

<b>Antibody Immunogen Source:</b>	Human PKCa (PRKCA) sequence peptide Cat. No.: PE-04AOU99
<b>Antibody Immunogen Sequence:</b>	SDP(pY)VKL(bA)C (bA) = beta-alanine
<b>Location in Target:</b>	Corresponds to amino acid residues S192 to L198; In the C2 domain. This is the major in vivo site of phosphorylation of PKCa.
<b>Peptide Type:</b>	For phosphosite-specific recognition of target.
<b>Target Phosphosite:</b>	Tyr-195

### Production

<b>Antibody Host Species:</b>	Rabbit
<b>Antibody Type:</b>	Polyclonal
<b>Antibody Ig Isotype Clone Lot:</b>	Immunoglobulin G
<b>Production Method:</b>	The immunizing peptide was produced by solid phase synthesis on a multipепptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. This antibody was also subject to negative purification over phosphotyrosine-agarose.
<b>Antibody Amount:</b>	25 µg
<b>Antibody Concentration:</b>	0.5 / 1.0 mg/ml
<b>Lot Number:</b>	150305 / 160121
<b>Storage Buffer:</b>	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol
<b>Storage Conditions and Stability:</b>	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.

# AB-PK764

## PKCa-pY195 Antibody



# KINEXUS

Address: 8755 Ash Street, Suite 1  
Vancouver, British Columbia,  
Canada V6P 6T3

Email: [info@kinexus.ca](mailto:info@kinexus.ca)  
Phone: 604-323-2547

### Applications

<b>Product Use:</b>	Western blotting   Antibody microarrays
<b>Antibody Dilution Recommended:</b>	2 µg/ml for immunoblotting
<b>Antibody Species Reactivity:</b>	Human, mouse, rat and many other mammals
<b>Antibody Positive Controls:</b>	Very strong immunoreactivity with immunogen peptide on dot blots.
<b>Overall Antibody Specificity:</b>	Very high selectivity
<b>Antibody Cross Reactivities:</b>	No significant cross-reactive proteins detected in phenylarsine oxide (PAO)+vanadate-treated HeLa cells, EGF-treated A431 cells and insulin-treated MCF7 cells, when these cells were homogenized in SDS-PAGE sample buffer.

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit [www.kinexusproducts.ca](http://www.kinexusproducts.ca) or contact us at 1-866-KINEXUS(546-3987)