## **AB-PK801**

### Ron-pY1238 +pY1239 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human protein-tyrosine kinase Ron (RONa)



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

 a r	$\sim$	Δt	 ra	tΔI	ın
a١	u	てに	ΙU	tei	
	J				

Name Long:	Macrophage-stimulating protein receptor alpha chain
Alias:	CD136; CD136 antigen; CDW136; C-met-related tyrosine kinase; Kinase RON; Macrophage stimulating 1 receptor; Macrophage-stimulating protein receptor precursor; MSP receptor; MST1R; PTK8; STK; CCDS2807.1; ENSG00000164078
UniProt ID:	Q04912
Sequence Predicted Mass (KDa):	152.271 (1400 AA; Q04912); 147.277 (1351 AA; Q04912-2); 97.315 (907 AA; Q04912-5); 69.265 (647 AA; Q04912-6); 57.721 (541 AA; Q04912-4); 52.519 (495 AA; Q04912-3)
Observed SDS-PAGE Mass (KDa):	150-220

#### Immunogen

Antibody Immunogen Source:	Human Ron (RONa) sequence peptide Cat. No.: PE-04ADM99		
Antibody Immunogen Sequence:	DRE(pY)(pY)SV(bA)C (bA) = beta-alanine		
Location in Target:	Corresponds to amino acid residues D1235 to V1241; In protein kinase catalytic domain activation T-loop between subdomains VII and VIII.		
Peptide Type:	For phosphosite-specific recognition of target.		
Target Phosphosite:	Tyr-1238 + Tyr-1239		

### Production

Antibody Host Species:	Rabbit		
Antibody Type:	Polyclonal		
Antibody Ig Isotype Clone Lot:	Immunoglobulin G		
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification over phosphotyrosine-agarose.		
Antibody Amount:	25 μg		
Antibody Concentration:	1 mg/ml		
Lot Number:	150106		
Storage Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol		
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.		

# AB-PK801 Ron-pY1238 +pY1239 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

### Applications

Product Use:	Western blotting   Antibody microarrays
Antibody Dilution Recommended:	2 μg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Antibody Positive Controls:	Very strong immunoreactivity with immunogen peptide on dot blots.
Overall Antibody Specificity:	Very high selectivity
<b>Antibody Cross Reactivities:</b>	No significant cross-reactive proteins detected in HeLa, HepG2 and Jurkat cells.

This product is for in vitro research use only and is not intended for use in humans or animals.