## AB-PK816 SLK-pS189 Antibody

SLK-pS189 Antibody Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human protein-serine/threonine kinase SLK



Email: info@kinexus.ca Phone: 604-323-2547

Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Target Protein	
Name Long:	STE20-like serine/threonine-protein kinase; CTCL tumour antigen se20-9
Alias:	bA16H23.1; KIAA0204; LOSK; MGC133067; Se20-9; STE20-like kinase; STK2
UniProt ID:	Q9H2G2
Sequence Predicted Mass (KDa):	142.695 (1235 AA; Q9H2G2); 138.996 (1204 AA; Q9H2G2-2)
Observed SDS-PAGE Mass (KDa):	150-200
Immunogen	
Antibody Immunogen Source:	Human SLK sequence peptide Cat. No.: PE-04ALC99
Antibody Immunogen Sequence:	RRD(pS)FIG(bA)C (bA) = beta-alanine
Location in Target:	Corresponds to amino acid residues R186 to G192; In the protein kinase catalytic domain activation T loop region between subdomains VII and VIII.
Peptide Type:	For phosphosite-specific recognition of target.
Target Phosphosite:	Ser-189
Production	
Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH
Antibody Amount:	<ul> <li>7.0 with saturated Tris. This antibody was also subject to negative purification over phosphotyrosine-agarose.</li> <li>25 µg</li> </ul>
Antibody Concentration:	0.75 mg/ml
Lot Number:	141202
Storage Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.

## AB-PK816 SLK-pS189 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications	
Product Use:	Western blotting   Antibody microarrays
Antibody Dilution Recommended:	2 µg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Antibody Positive Controls:	Very strong immunoreactivity with immunogen peptide on dot blots.
Detection by Immunoblotting in Cell/Tissue Lysates:	Weak immunoreactivity of a target-sized protein by Western blotting in MCF7 cells.
Overall Antibody Specificity:	High-very high selectivity
Antibody Cross Reactivities:	No significant cross-reactive proteins detected in HEK-293, HeLa and Jurkat cells, except a 50 KDa protein in A431 cells and 250 KDa protein in HEK-293 and MCF7 cells.

This product is for in vitro research use only and is not intended for use in humans or animals.