AB-PK827 TAO1-pY309 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human protein-serine/threonine kinase TAO1 (TAOK1)

Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3



Email: info@kinexus.ca Phone: 604-323-2547

	1 -		
loro	Ot L	roti	מור
Targ			
	O	100	2111

Name Long:	Serine/threonine-protein kinase TAO1	
Alias:	FLJ14314; HKFC-B; KIAA1361; MAP3K16; MARKK; PSK2; STE20-like kinase PSK2; TAOK1; TAO kinase 1; KFC-B; MARKK; h KFC-B; ENSG00000160551	
UniProt ID:	Q7L7X3	
Sequence Predicted Mass (KDa):	116.070 (1001 AA; Q7L7X3); 97.526 (853 AA; Q7L7X3-3); 46.163 (398 AA; Q7L7X3-2)	
Observed SDS-PAGE Mass (KDa):	110-125	

Immunogen

Antibody Immunogen Source:	Human TAO1 (TAOK1) sequence peptide Cat. No.: PE-04APX95	
Antibody Immunogen Sequence:	NLQ(pY)RKM(bA)C (bA) = beta-alanine	
Location in Target:	Corresponds to amino acid residues N306 to M312; Just after the kinase catalytic domain. Thhis is the major in vivo site of phosphorylation on TAO1.	
Peptide Type:	For phosphosite-specific recognition of target.	
Target Phosphosite:	Tyr-309	

Production

Troduction		
Antibody Host Species:	Rabbit	
Antibody Type:	Polyclonal	
Antibody Ig Isotype Clone Lot:	Immunoglobulin G	
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. This antibody was also subject to negative purification over phosphotyrosine-agarose.	
Antibody Amount:	25 μg	
Antibody Concentration:	1 mg/ml	
Lot Number:	150305	
Storage Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol	
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.	

AB-PK827 TAO1-pY309 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications	
Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	2 μg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals; Phosphosite is highly conserved in mammals
Antibody Positive Controls:	Very strong immunoreactivity with immunogen peptide on dot blots.
Detection by Immunoblotting in Cell/Tissue Lysates:	Strong immunoreactivity of a target-sized protein by Western blotting in HeLa cells and weak detection in HepG2 cells.
Overall Antibody Specificity:	High selectivity
Antibody Cross Reactivities:	No significant cross-reactive proteins detected in HepG2, MCF7 and T98G cells, except for ~80 KDa protein in HeLa cells and ~65 KDa protein in inulin-treated MCF7 cells.