## **AB-PK848**

### Tyro3-pY685+pY686 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human protein-tyrosine kinase Tyro3



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

#### **Target Protein**

Name Long: Tyrosine-protein kinase receptor TYRO3

**Alias:** BYK; DTK; RSE; SKY; Tif; Brt; FLJ16467; CCDS10080.1; ENSG00000092445

UniProt ID: Q06418

Sequence Predicted Mass (KDa): 96.905 (890 AA; Q06418)

Observed SDS-PAGE Mass (KDa): 95-110

#### Immunogen

Antibody Immunogen Source:	Human Tyro3 sequence peptide Cat. No.: PE-04AQL90
Antibody Immunogen Sequence:	SGD(pY)(pY)RQG(bA)C (bA) = beta-alanine
Location in Target:	Corresponds to amino acid residues S682 to G689; In the protein kinase catalytic domain activation T loop region between subdomains VII and VIII.
Peptide Type:	For phosphosite-specific recognition of target.
Target Phosphosite:	Tyr-685+Tyr-686

#### Production

Antibody Type:  Antibody Ig Isotype Clone Lot:	Rabbit Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep
Antibody Ig Isotype Clone Lot:	Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep
Ţ	The immunizing peptide was produced by solid phase synthesis on a multipep
ļ.	
Production Method:  i  i  i  i  i  i  i  i  i  i  i  i  i	peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. This antibody was also subject to negative purification over phosphotyrosine-agarose.
Antibody Amount:	25 μg
Antibody Concentration:	1 mg/ml
Lot Number:	150305
Storage Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol
	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.

# AB-PK848 Tyro3-pY685+pY686 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

### Applications

Product Use:	Western blotting   Antibody microarrays
Antibody Dilution Recommended:	2 μg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Antibody Positive Controls:	Very strong immunoreactivity with immunogen peptide on dot blots.
Detection by Immunoblotting in Cell/Tissue Lysates:	Medium immunoreactivity of a target-sized protein by Western blotting in HepG2 cells.
Overall Antibody Specificity:	High selectivity
Antibody Cross Reactivities:	No significant cross-reactive proteins detected in MCF7 cells, except for a strong ~140 KDa protein, which was also detected in HepG2 cells. ~75 and 85 KDa proteins that were also detectable in HepG2 cells.

This product is for in vitro research use only and is not intended for use in humans or animals.