AB-PK861 ZAP70-pY292 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human protein-tyrosine kinase ZAP70



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

| | 1 | | 1 - ' - |
|------|------|-----|---------|
| I Or | пот. | urα | TOID |
| | | Pro | |
| | 900 | | |

| Name Long: | Zeta-chain (TCR) associated protein-tyrosine kinase, 70 kDa |
|--------------------------------|---|
| Alias: | SRK; Syk-related tyrosine kinase; ZA70; ZAP-70 |
| UniProt ID: | P43403 |
| Sequence Predicted Mass (KDa): | 69.872 (619 AA; P43403); 55.873 (493 AA; P43403-3); 35.647 (312 AA; P43403-2) |
| Observed SDS-PAGE Mass (KDa): | 65-75 |

Immunogen

| Antibody Immunogen Source: | Human ZAP70 sequence peptide Cat. No.: PE-04AGT85 | |
|-------------------------------------|---|--|
| Antibody Immunogen Sequence: | SDG(pY)TPE(bA)C (bA) = beta-alanine | |
| Location in Target: | Corresponds to amino acid residues S289 to E295; In the region between the SH2 domain and the protein kinase catalytic domain | |
| Peptide Type: | For phosphosite-specific recognition of target. | |
| Target Phosphosite: | Tyr-292 | |

Production

| Antibody Type: Polyclonal Antibody Ig Isotype Clone Lot: Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. This antibody was also subject to negative purification over phosphotyrosine-agarose. Antibody Amount: 25 μg Antibody Concentration: 1 mg/ml Lot Number: 150205 Storage Buffer: Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol Storage Conditions and Stability: For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles. | | | |
|--|-----------------------------------|--|--|
| Antibody Ig Isotype Clone Lot: Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. This antibody was also subject to negative purification over phosphotyrosine-agarose. Antibody Amount: 25 µg Antibody Concentration: 1 mg/ml Lot Number: 150205 Storage Buffer: Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol For long term storage, keep frozen at -40°C or lower. Stock solution can be kept | Antibody Host Species: | Rabbit | |
| The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. This antibody was also subject to negative purification over phosphotyrosine-agarose. Antibody Amount: 25 µg Antibody Concentration: 1 mg/ml Lot Number: 150205 Storage Buffer: Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol For long term storage, keep frozen at -40°C or lower. Stock solution can be kept | Antibody Type: | Polyclonal | |
| peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification over phosphotyrosine-agarose. Antibody Amount: 25 μg Antibody Concentration: 1 mg/ml Lot Number: 150205 Storage Buffer: Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol For long term storage, keep frozen at -40°C or lower. Stock solution can be kept | Antibody Ig Isotype Clone Lot: | Immunoglobulin G | |
| Antibody Concentration: 1 mg/ml Lot Number: 150205 Storage Buffer: Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol For long term storage, keep frozen at -40°C or lower. Stock solution can be kept | Production Method: | peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification | |
| Lot Number: 150205 Storage Buffer: Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol Storage Conditions and Stability: For long term storage, keep frozen at -40°C or lower. Stock solution can be kept | Antibody Amount: | 25 μg | |
| Storage Buffer: Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol For long term storage, keep frozen at -40°C or lower. Stock solution can be kept | Antibody Concentration: | 1 mg/ml | |
| Storage Conditions and Stability: For long term storage, keep frozen at -40°C or lower. Stock solution can be kept | Lot Number: | 150205 | |
| | Storage Buffer: | Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol | |
| | Storage Conditions and Stability: | | |

AB-PK861 ZAP70-pY292 Antibody

Applications



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

| Applications | | |
|---|---|--|
| Product Use: | Western blotting Antibody microarrays | |
| Antibody Dilution Recommended: | 2 μg/ml for immunoblotting | |
| Antibody Species Reactivity: | Human, mouse, rat and many other mammals | |
| Detection by Immunoblotting in Cell/Tissue Lysates: | Weak immunoreactivity of a target-sized protein by Western blotting in A431 and Jurkat cells. | |
| Overall Antibody Specificity: | High selectivity | |
| Antibody Cross Reactivities: | Almost no significant cross-reactivities detected inMCF7 cells, except for 80-130 KDa cross-reactive proteins detected with SDS-PAGE sample buffer extracted MCF7 cells. No cross-reactivities were seen in A431 cells (except strong 200 | |

KDa protein cross-reactivity), Jurkat and HeLa cells.

This product is for in vitro research use only and is not intended for use in humans or animals.