# AB-PK863 ZAP70-pY492+pY493 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human protein-tyrosine kinase ZAP70



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

	1	
loro	Jrat	'AID
Targ		еш

Name Long:	Zeta-chain (TCR) associated protein-tyrosine kinase, 70 kDa
Alias:	SRK; Syk-related tyrosine kinase; ZA70; ZAP-70
UniProt ID:	P43403
Sequence Predicted Mass (KDa):	69.872 (619 AA; P43403); 55.873 (493 AA; P43403-3); 35.647 (312 AA; P43403-2)
Observed SDS-PAGE Mass (KDa):	65-75

#### Immunogen

Antibody Immunogen Source:	Human ZAP70 sequence peptide Cat. No.: PE-04AGS80
Antibody Immunogen Sequence:	DDS(pY)(pY)TAR(bA)C (bA) = beta-alanine
Location in Target:	Corresponds to amino acid residues D489 to R496; In the protein kinase catalytic domain.
Peptide Type:	For phosphosite-specific recognition of target.
Target Phosphosite:	Tyr-492+Tyr-493

#### Production

Antibody Host Species:	Rabbit	
Antibody Type:	Polyclonal	
Antibody Ig Isotype Clone Lot:	Immunoglobulin G	
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification over phosphotyrosine-agarose.	
Antibody Amount:	25 μg	
Antibody Concentration:	1 mg/ml	
Lot Number:	150205	
Storage Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol	
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.	

## AB-PK863 ZAP70-pY492+pY493 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

### Applications

Product Use:	Western blotting   Antibody microarrays
Antibody Dilution Recommended:	2 μg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Detection by Immunoblotting in Cell/Tissue Lysates:	Weak immunoreactivity of a target-sized protein by Western blotting in HeLa and Jurkat cells treated with vanadate and phenylarsine oxide for 30 minutes.
Overall Antibody Specificity:	High selectivity
Antibody Cross Reactivities:	Almost no significant cross-reactivities detected in HeLa, Jurkat and MCF7 cells, except phenylarsine oxide (PAO) increases 70, 50 + 34 KDa cross-reactive proteins in Jurkat and HeLa cells.

This product is for in vitro research use only and is not intended for use in humans or animals.