AB-PN510 EML4-pY226 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human EML4



Email: info@kinexus.ca Phone: 604-323-2547

Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Name Long:	Echinoderm microtubule-associated protein-like 4
Alias:	C2orf2; DKFZp686P18118; echinoderm microtubule associated protein like 4; Echinoderm microtubule-associated protein-like 4; ELP120; EMAL4; EMAP-4; EMAPL4; EML4; FLJ10942; FLJ32318; Restrictedly overexpressed proliferation- associated protein; Ropp 120; ROPP120
UniProt ID:	Q9HC35
Sequence Predicted Mass (KDa):	108.916 (981 AA; Q9HC35-1); 102.444 (923 AA; Q9HC35-2)
Observed SDS-PAGE Mass (KDa):	115-125
Immunogen	
Antibody Immunogen Source:	Human EML4 sequence peptide Cat. No.: PE-04AHG99
Antibody Immunogen Sequence:	EGE(pY)IKM(bA)C (bA) = beta-alanine
Location in Target:	Corresponds to amino acid residues E223 to M229; In the N-terminal portion of EML4 near the HELP domain. This is the major in vivo phosphorylation site in EML4.
Peptide Type:	For phosphosite-specific recognition of target.
Target Phosphosite:	Tyr-226
Production	Debbit
Antibody Host Species:	Rabbit
	Polyclonal
Antibody Host Species: Antibody Type: Antibody Ig Isotype Clone Lot:	
Antibody Type: Antibody Ig Isotype Clone Lot:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with
Antibody Type: Antibody Ig Isotype Clone Lot: Production Method:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification
Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: Antibody Amount:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to ph 7.0 with saturated Tris. This antibody was also subject to negative purification over phosphotyrosine-agarose.
Antibody Type: Antibody Ig Isotype Clone Lot: Production Method: Antibody Amount: Antibody Concentration:	Polyclonal Immunoglobulin G The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification over phosphotyrosine-agarose. 25 μg
Antibody Type:	PolyclonalImmunoglobulin GThe immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. This antibody was also subject to negative purification over phosphotyrosine-agarose.25 μg1 mg/ml

AB-PN510 EML4-pY226 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications	
Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	2 µg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Antibody Positive Controls:	Very strong immunoreactivity with immunogen peptide on dot blots.

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit <u>www.kinexusproducts.ca</u> or contact us at 1-866-KINEXUS(546-3987)