# AB-PN518 HCA59-pY147 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human HCA59 (Q9NZ63)

KINEXUS

Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia,

Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

## Target Protein

Name Long:	Uncharacterized protein C9orf78
Alias:	bA409K20.3; C9orf78; chromosome 9 open reading frame 78; Cl078; HCA59; Hepatocellular carcinoma-associated antigen 59; HSPC220
UniProt ID:	Q9NZ63
Sequence Predicted Mass (KDa):	33.688 (289 AA; Q9NZ63)
Observed SDS-PAGE Mass (KDa):	35-39

#### Immunogen

Antibody Immunogen Source:	Human HCA59 (Q9NZ63) sequence peptide Cat. No.: PE-04ACZ80
Antibody Immunogen Sequence:	DCL(pY)ELP(bA)C (bA) = beta-alanine
Location in Target:	Corresponds to amino acid residues D144 to P150; In the middle of the Hep_59 domain. This is the major in vivo phosphorylation site in HCA59.
Peptide Type:	For phosphosite-specific recognition of target.
Target Phosphosite:	Tyr-147

### Production

Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification over phosphotyrosine-agarose.
Antibody Amount:	25 μg
Antibody Concentration:	1 mg/ml
Lot Number:	150106
Storage Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.

# AB-PN518 HCA59-pY147 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

### Applications

Product Use:	Western blotting   Antibody microarrays
Antibody Dilution Recommended:	2 μg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Antibody Positive Controls:	Very strong immunoreactivity with immunogen peptide on dot blots.
Overall Antibody Specificity:	Very high selectivity
Antibody Cross Reactivities:	No significant cross-reactive proteins detected in A431 cells.

This product is for in vitro research use only and is not intended for use in humans or animals.