AB-PN523 PECAM-1-pY713 Antibody

Phosphosite-specific polyclonal antibody for monitoring the phosphorylation of human PECAM-1



Email: info@kinexus.ca Phone: 604-323-2547

Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Target Protein	
Name Long:	Platelet endothelial cell adhesion molecule
Alias:	Adhesion molecule; CD31; CD31 antigen; CD31/EndoCAM; EndoCAM; FLJ58394; GPIIA'; PECA1; PECAM-1; PECAM-1, CD31/EndoCAM; PECAM1; Platelet endothelial cell adhesion molecule; platelet/endothelial cell adhesion molecule
UniProt ID:	P16284
Sequence Predicted Mass (KDa):	82.536 (738 AA; P16284-1); 81.378 (727 AA; P16284-6); 80.479 (720 AA; P16284-2); 80.419 (719 AA; P16284-4); 80.207 (717 AA; P16284-3); 79.261 (708 AA; P16284-5)
Observed SDS-PAGE Mass (KDa):	125-135
Immunogen	
Antibody Immunogen Source:	Human PECAM-1 sequence peptide Cat. No.: PE-04AEZ99
Antibody Immunogen Sequence:	ETV(pY)SEV(bA)C (bA) = beta-alanine
Location in Target:	Corresponds to amino acid residues E710 to V716; In the C-terminal region of the protein after the transmembrane domain. This is the major in vivo phosphorylation site in PECAM-1.
Peptide Type:	For phosphosite-specific recognition of target.
Target Phosphosite:	Tyr-713
Production Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Ig Isotype Clone Lot:	Immunoglobulin G
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from each animal was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.This antibody was also subject to negative purification over phosphotyrosine-agarose.
Antibody Amount:	25 μg
Antibody Concentration:	1 mg/ml
Lot Number:	150203
Storage Buffer:	Phosphate buffered saline (PBS) pH7.4, 0.05% Thimerasol
Storage Conditions and Stability:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.

AB-PN523 PECAM-1-pY713 Antibody



Address: 8755 Ash Street, Suite 1 Vancouver, British Columbia, Canada V6P 6T3

Email: info@kinexus.ca Phone: 604-323-2547

Applications	
Product Use:	Western blotting Antibody microarrays
Antibody Dilution Recommended:	2 µg/ml for immunoblotting
Antibody Species Reactivity:	Human, mouse, rat and many other mammals
Overall Antibody Specificity:	Medium-High selectivity
Antibody Cross Reactivities:	In Jurkat cells, phenylarsine oxide (PAO) increases detection of 75-85 KDa and 15 KDa cross-reactive proteins.

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit <u>www.kinexusproducts.ca</u> or contact us at 1-866-KINEXUS(546-3987)