

AB-PN640

p53-pS392 Antibody

Phosphosite-specific rabbit polyclonal antibody for TP53 (p53). This phosphoserine-site antibody is highly specific for phosphoserine in the phosphosite.

Website: www.kinexus.ca
Email: info@kinexus.ca
Phone: 604-323-2547



Address: 8755 Ash Street, Suite 1
Vancouver, British Columbia,
Canada V6P 6T3

Target Protein

Protein Name Long:	Tumour suppressor protein p53 (antigenNY-CO-13)
Protein Alias:	P53; LFS1; TRP53; FLJ92943; TP53; p53 Tumour Suppressor; Phosphoprotein p53; TP53; Transformation related protein 53; TRP53; Tumour protein p53
UniProt ID:	P04637
Protein Molecular Mass:	43,653 Da (393 AA; P04637-1); 39,320 Da (354 AA; P04637-4); 37,826 Da (341 AA; P04637-2); 34,168 Da (307 AA; P04637-6); 29,553 Da (261 AA; P04637-7); 23,726 Da (209 AA; P04637-8)

Immunogen

Antibody Immunogen Source:	Synthetic phosphopeptide patterned after human p53
Antibody Immunogen Sequence:	CEGPD(pS)D(-cooh)
Antibody Immunogen Description:	Corresponds to amino acid residues E388 to D393. S392 phosphorylation induces activity, stimulates binding to DNA, NFkB-p65, UBE4B and p21Cip1, and inhibits binding of CDK1 and MDM2. This is one of the minor <i>in vivo</i> phosphorylation sites in p53 (≥ 13 reports by mass spectrometry in PhosphositePlus), although there are ≥ 94 reports with Western blotting. TP53 is known to be phosphorylated at this site <i>in vitro</i> by ATM, CDK7, CDK9, CK2a1 (CSNK2A1), ERK1 (MAPK3), LKB1 (STK11), NuaK1, and PKR (EIF2AK2).
Antibody Target Type:	Phosphosite-specific

Production

Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Isotype:	IgG
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multi-peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from these animals was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Amount:	25 μ g
Antibody Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline pH 7.4, 0.05% Thimerasol
Storage Conditions:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.
Storage Stability:	>2 years

Applications

Product Use:	Western blotting Antibody microarray
Antibody Dilution Recommended:	2 µg/ml for immunoblotting
Antibody Species Reactivity:	Human Chimpanzee Rhesus macaque Rat Mouse Frog Zebra fish

This product is for *in vitro* research use only and is not intended for use in humans or animals.

For more information on our products please visit <https://kinexus-ca.myshopify.com/> or contact us at 1-866-546-3987