

AB-PN716

PFKP-pS386 Antibody

Phosphosite-specific rabbit polyclonal antibody for PFKP. This phosphoserine-site antibody also cross-reacts with phosphothreonine and weakly with a phosphotyrosine substitution in the phosphosite.

Website: www.kinexus.ca

Email: info@kinexus.ca

Phone: 604-323-2547



Address: 8755 Ash Street, Suite 1
Vancouver, British Columbia,
Canada V6P 6T3

Target Protein

Protein Name Long:	6-phosphofructokinase type C
Protein Alias:	6-phosphofructokinase type C; Phosphofructokinase 1; Phosphohexokinase; Phosphofructo-1-kinase isozyme C; 6-phosphofructokinase, platelet type; Phosphofructokinase-P; Phosphofructokinase, platelet; 6-phosphofructokinase, platelet type; K6PP; PFK, platelet type; PFK-C; PFKF; Phosphofructo-1-kinase isozyme C; Phosphofructokinase 1; Phosphofructokinase, platelet; Phosphofructokinase, platelet type; Phosphohexokinase
UniProt ID:	Q01813
Protein Molecular Mass:	85,596 Da (784 AA)

Immunogen

Antibody Immunogen Source:	Synthetic phosphopeptide patterned after human PFKP
Antibody Immunogen Sequence:	RGR(pS)FAG(β A)C
Antibody Immunogen Description:	Corresponds to amino acid residues R383 to G389. The effect of S386 phosphorylation is unclear. This is the major <i>in vivo</i> phosphorylation site in PFKP (≥ 234 reports from high throughput mass spectrometry studies recorded in PhosphoSitePlus). This human phosphosite is highly conserved in vertebrates and also found in the fruit fly.
Antibody Target Type:	Phosphosite-specific

Production

Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Isotype:	IgG
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from these animals was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Amount:	25 μ g
Antibody Concentration:	1 mg/ml
Storage Buffer:	Phosphate buffered saline pH 7.4, 0.05% Thimerasol
Storage Conditions:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at $+4^{\circ}\text{C}$ for more than 3 months. Avoid repeated freeze-thaw cycles.
Storage Stability:	>2 years

Applications

Product Use:	Western blotting Antibody microarray
Antibody Dilution Recommended:	2 µg/ml for immunoblotting
Antibody Species Reactivity:	Human Dog Rat Platypus Chicken Frog Zebra fish Fruit fly

This product is for *in vitro* research use only and is not intended for use in humans or animals.

For more information on our products please visit <https://kinexus-ca.myshopify.com/> or contact us at 1-866-546-3987