

AB-PN983

RTN4-pS107 Antibody

Phosphosite-specific rabbit polyclonal antibody for RTN4 (NOGO). This phosphoserine-site antibody also cross-reacts with phosphothreonine, phosphotyrosine, serine, threonine and glutamic acid substitutions in the

Website: www.kinexus.ca
Email: info@kinexus.ca
Phone: 604-323-2547



Address: 8755 Ash Street, Suite 1
Vancouver, British Columbia,
Canada V6P 6T3

Target Protein

Protein Name Long:	Reticulon-4
Protein Alias:	ASY; Foccen; KIAA0886; Neuroendocrine-specific protein; Neuroendocrine-specific protein C homolog; NOGO; Nogo protein; NSP; NSP-CL; Reticulon 4; Reticulon-4; Reticulon-5; RTN-x
UniProt ID:	Q9NQC3
Protein Molecular Mass:	129,931 Da (1,192 AA)

Immunogen

Antibody Immunogen Source:	Synthetic phosphopeptide patterned after human RTN4
Antibody Immunogen Sequence:	C(βA)ERQP(pS)WDP
Antibody Immunogen Description:	Corresponds to amino acid residues E103 to P110. The effect of Nogo S107 phosphorylation is unclear. This is the major site of <i>in vivo</i> phosphorylation of RTN4 (Nogo) based on ≥369 mass spectrometry reports recorded in PhosphoSitePlus. RTN4 is known to be phosphorylated at this site <i>in vitro</i> by MAPKAPK2.
Antibody Target Type:	Phosphosite-specific

Production

Antibody Host Species:	Rabbit
Antibody Type:	Polyclonal
Antibody Isotype:	IgG
Production Method:	The immunizing peptide was produced by solid phase synthesis on a multipep peptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from these animals was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris.
Amount:	25 µg
Antibody Concentration:	0.5 mg/ml
Storage Buffer:	Phosphate buffered saline pH 7.4, 0.05% Thimerasol
Storage Conditions:	For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles.
Storage Stability:	>2 years

Applications

Product Use:	Western blotting Antibody microarray
Antibody Dilution Recommended:	2 µg/ml for immunoblotting
Antibody Species Reactivity:	Human Rhesus macaque Rat Mouse

This product is for *in vitro* research use only and is not intended for use in humans or animals.

For more information on our products please visit <https://kinexus-ca.myshopify.com/> or contact us at 1-866-546-3987