

AB-PP556

PTEN-pY336 Antibody

Phosphosite-specific rabbit polyclonal antibody for PTEN. This phosphotyrosine-site antibody is highly specific for phosphotyrosine.

Website: www.kinexus.ca
Email: info@kinexus.ca
Phone: 604-323-2547



Address: 8755 Ash Street, Suite 1
Vancouver, British Columbia,
Canada V6P 6T3

Target Protein

| | |
|--------------------------------|---|
| Protein Name Long: | Phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase and protein phosphatase and tensin homologue deleted on chromosome 10 |
| Protein Alias: | BZS; MHAM; MMAC1; Mutated in multiple advanced cancers 1; Phosphatase and tensin; Phosphatase and tensin homolog gene; Protein-tyrosine phosphatase PTEN; PTEN1; TEP1 |
| UniProt ID: | P60484 |
| Protein Molecular Mass: | 47,166 Da (403 AA; P60484-1); 64,882 Da (576 AA; P60484-2); 19,796 Da (175 AA; P60484-3) |

Immunogen

| | |
|--|---|
| Antibody Immunogen Source: | Synthetic phosphopeptide patterned after human PTEN |
| Antibody Immunogen Sequence: | KANR(pY)FSP(βA)C |
| Antibody Immunogen Description: | Corresponds to amino acid residues K332 to P339. Y336 phosphorylation stimulates PTEN phosphatase activity and protein stability, which in turn prevents its ubiquitination and degradation, possibly by reducing its binding to NEDD4. Frk binds (via its SH3-domain) with PTEN. This site was not identified as a site of <i>in vivo</i> phosphorylation of PTEN based mass spectrometry reports recorded in PhosphoSitePlus. PTEN is known to be phosphorylated at this site <i>in vitro</i> by Frk. |
| Antibody Target Type: | Phosphosite-specific |

Production

| | |
|--------------------------------|---|
| Antibody Host Species: | Rabbit |
| Antibody Type: | Polyclonal |
| Antibody Isotype: | IgG |
| Production Method: | The immunizing peptide was produced by solid phase synthesis on a multipепptide synthesizer and purified by reverse-phase hplc chromatography. Purity was assessed by analytical hplc and the amino acid sequence confirmed by mass spectrometry analysis. This peptide was coupled to KLH prior to immunization into rabbits. New Zealand White rabbits were subcutaneously injected with KLH-coupled immunizing peptide every 4 weeks for 4 months. The sera from these animals was applied onto an agarose column to which the immunogen peptide was thio-linked. Antibody was eluted from the column with 0.1 M glycine, pH 2.5. Subsequently, the antibody solution was neutralized to pH 7.0 with saturated Tris. |
| Amount: | 25 µg |
| Antibody Concentration: | 0.35 mg/ml |
| Storage Buffer: | Phosphate buffered saline pH 7.4, 0.05% Thimerasol |
| Storage Conditions: | For long term storage, keep frozen at -40°C or lower. Stock solution can be kept at +4°C for more than 3 months. Avoid repeated freeze-thaw cycles. |
| Storage Stability: | >2 years |

Applications

| | |
|---------------------------------------|---|
| Product Use: | Western blotting Antibody microarray |
| Antibody Dilution Recommended: | 2 µg/ml for immunoblotting |
| Antibody Species Reactivity: | Human Rhesus macaque Rat Mouse Platypus |

This product is for *in vitro* research use only and is not intended for use in humans or animals.

For more information on our products please visit <https://kinexus-ca.myshopify.com/> or contact us at 1-866-546-3987