

LC0205S

HEK293+Forskolin Cytosol

Cytosolic lysate boiled in SDS-PAGE sample buffer prepared from HEK293 cells that were cultivated to 90% confluency and deprived of serum for 18 hours and then treated with 150 nM forskolin for 15 minutes prior to harvesting

Address: 8755 Ash Street, Suite 1
Vancouver, British Columbia,
Canada V6P 6T3



Email: info@kinexus.ca
Phone: 604-323-2547

Production

Product Name Long:	Human Embryonic Kidney-293 cells - Forskolin-treated - Cytosolic lysate
Production Method:	HEK293 cells were cultivated to 80% confluency and deprived of serum for 18 hours and then treated with 150 nM forskolin for 15 minutes prior to harvesting. Lysates were prepared from scrapped cells that were homogenized by sonication in buffer formulated with 60 mM β -glycerophosphate, pH 7.2, 20 mM MOPS, 20 mM sodium pyrophosphate, 30 mM sodium fluoride, 5 mM EDTA, 3 mM benzamidine, 2 mM EGTA, 1 mM sodium orthovanadate, 1 mM phenylmethylsulfonylfluoride, 1 mM dithiothreitol, 10 μ M leupeptin, and 5 μ M pepstatin A. Cytosolic lysates were prepared following sonication and 30 min ultracentrifugation at 100,000 rpm. Lysates were further diluted in SDS-PAGE sample buffer at a final concentration of 2 mg/ml.
Amount:	200 μ g
Protein Concentration:	2 mg/ml
Storage Stability:	1 year at -70°C

Applications

Lysate Use Description:	For testing antibodies by immunoblotting.
--------------------------------	---

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit www.kinexusproducts.ca or contact us at 1-888-546-3987