

LC0504S

HL-60+FBS Cytosol

Cytosolic lysate boiled in SDS-PAGE sample buffer prepared from HL-60 cells that were cultivated to 80% confluency in the presence of 20% fetal bovine serum prior to harvesting



KINEXUS

Address: 8755 Ash Street, Suite 1
Vancouver, British Columbia,
Canada V6P 6T3

Email: info@kinexus.ca
Phone: 604-323-2547

Production

Product Name Long:	Human promyelocytic HL-60 cells - Cytosolic lysate
Production Method:	HL-60 cells were cultivated to 80% confluency in the presence of 20% fetal bovine serum prior to harvesting. Lysates were prepared from scrapped cells that were homogenized by sonication in buffer formulated with 60 mM β -glycerophosphate, pH 7.2, 20 mM MOPS, 20 mM sodium pyrophosphate, 30 mM sodium fluoride, 5 mM EDTA, 3 mM benzamidine, 2 mM EGTA, 1 mM sodium orthovanadate, 1 mM phenylmethylsulfonylfluoride, 1 mM dithiothreitol, 10 μ M leupeptin, and 5 μ M pepstatin A. Cytosolic lysates were prepared following sonication and 30 min ultracentrifugation at 100,000 rpm. Lysates were further diluted in SDS-PAGE sample buffer at a final concentration of 2 mg/ml.
Amount:	200 μ g
Protein Concentration:	2 mg/ml
Storage Stability:	1 year at -70°C

Applications

Lysate Use Description:	For testing antibodies by immunoblotting.
--------------------------------	---

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit www.kinexusproducts.ca or contact us at 1-888-546-3987