

LC0514F

HL-60+Staurosp. Cytosol

Cytosolic lysate from HL-60 cells that were cultivated to 80% confluency in the presence of 20% fetal bovine serum and then treated with 0.5 µg/ml staurosporine for 15 minutes prior to harvesting



KINEXUS

Address: 8755 Ash Street, Suite 1
Vancouver, British Columbia,
Canada V6P 6T3

Email: info@kinexus.ca
Phone: 604-323-2547

Production

Product Name Long:	Human promyelocytic HL-60 cells - Staurosporine-treated - Cytosolic lysate
Production Method:	HL-60 cells were cultivated to 80% confluency in the presence of 20% fetal bovine serum and then treated with 0.5 µg/ml staurosporine for 15 minutes prior to harvesting. Lysates were prepared from scrapped cells that were homogenized by sonication in buffer formulated with 60 mM β-glycerophosphate, pH 7.2, 20 mM MOPS, 20 mM sodium pyrophosphate, 30 mM sodium fluoride, 5 mM EDTA, 3 mM benzamidine, 2 mM EGTA, 1 mM sodium orthovanadate, 1 mM phenylmethylsulfonylfluoride, 1 mM dithiothreitol, 10 µM leupeptin, and 5 µM pepstatin A. Cytosolic lysates were prepared following sonication and 30 min ultracentrifugation at 100,000 rpm. Lysates were further diluted in homogenizing buffer at a final concentration of 3 mg/ml.
Amount:	200 µg
Protein Concentration:	3 mg/ml
Storage Stability:	1 year at -70°C

Applications

Lysate Use Description:	For testing antibodies by immunoprecipitation or immunoblotting, and for assays of enzymes.
--------------------------------	---

This product is for in vitro research use only and is not intended for use in humans or animals.

For more information on our products please visit www.kinexusproducts.ca or contact us at 1-888-546-3987